

D5-46  
79. 30  
9186

N86 - 28540

STRONTIUM AND NEODYMIUM ISOTOPIC VARIATIONS IN EARLY ARCHEAN GNEISSES  
AFFECTED BY MIDDLE TO LATE ARCHEAN HIGH-GRADE METAMORPHIC PROCESSES: WEST  
GREENLAND AND LABRADOR. K.D. Collerson<sup>1,5</sup>, M.T. McCulloch<sup>1</sup>, D. Bridgwater<sup>2</sup>,  
V.R. McGregor<sup>3</sup> and A.P. Nutman<sup>4</sup>

<sup>1</sup> Research School of Earth Sciences, Australian National University, Canberra,  
Australia. <sup>2</sup> Geological Museum, Copenhagen, Denmark. <sup>3</sup> ~~Narsarsuaq~~ <sup>(A)</sup> ~~Greenland~~ <sup>(B)</sup>  
and <sup>4</sup> Geological Survey of Greenland, Copenhagen, Denmark, <sup>5</sup> Present Address  
Department of Geology, University of Regina, Regina, Saskatchewan, Canada.

Introduction. Relicts of continental crust formed more than 3400 Ma  
ago are preserved fortuitously in most cratons. They provide the most direct  
information about crust and mantle evolutionary processes during the first  
billion years of Earth history. In view of their polymetamorphic character,  
these terranes are commonly affected by subsequent tectonothermal events. Hence,  
their isotope systematics may be severely disturbed as a result of bulk chemical  
change or local isotopic homogenization. This leads to equivocal age and  
source information for different components within these terranes (see for  
example, [1-3]). In this paper we present Sr and Nd isotopic data for early  
Archean gneisses from the North Atlantic Craton in west Greenland and northern  
Labrador which have been affected by younger metamorphic events.

#### Regional Geological Setting.

(1) West Greenland. The early Archean Amitsoq gneisses of southern West  
Greenland were first identified around the mouth of Ameralik Fjord, ca. 15  
km south of Nuuk (Godthaab). In this area, both field and isotopic data document  
the existence of an ancient high-grade gneiss complex [4-6]. It is important  
to note that late Archean tectonothermal events (3100 to 2600 Ma) in this  
area were characterized by amphibolite facies rather than granulite facies  
metamorphic conditions, and that little migmatization occurred during this  
time interval. To the south and east, late Archean metamorphic grade reached  
granulite facies at 2900 - 2800 Ma, and mid-Archean tonalitic gneisses (correlated  
with 3050-2900 Ma Nuk gneisses in Godthaabsfjord) as well as later Archean  
granitoid gneisses, are volumetrically important. However, field studies  
suggest that an early component (Amitsoq gneiss) is present within these  
polyphase gneisses [7]. Relevant aspects of the regional geology are discussed  
in detail by McGregor *et al.*, [7] and Nutman *et al.*, [8]. Isotopic data for  
the gneisses are discussed in the context of three regional geologic field  
divisions:

(i) Amitsoq gneisses from type area (northwestern Buksefjorden and outer  
Ameralik). In this area there is evidence of ca. 3600 Ma granulite facies  
metamorphism [9].

ii) Amitsoq gneisses (from Nordafar) which are continuous with those in the  
type area but are strongly affected by late Archean granitoid injection and  
variable metasomatic alteration. The late granites probably formed as a result  
of granulite facies metamorphism at depth.

iii) Gneisses further to the south and east (at Kangimut sangmissoq, Tinissaq  
and Ikerasakisup akornga) which preserve field characteristics of Amitsoq  
gneisses but were metamorphosed under granulite facies conditions at ca. 2900  
to 2800 Ma and subsequently experienced patchy retrogression under amphibolite  
facies conditions. These units are not structurally continuous with the type  
Amitsoq gneisses to the northwest.


(2) Northern Labrador

The Uivak 1 gneisses and the interleaved Nulliak (supracrustal) assemblage in northern Labrador are of similar age and character to rocks described above from west Greenland [10-13]. Large areas of northern Labrador show variable effects of late Archean (ca. 2800 to 2900 Ma) granulite facies metamorphism. Early Archean relationships are best preserved on Big Island, Cape Uivak and the area immediately to the west and southwest, where late Archean metamorphic grade did not exceed amphibolite facies conditions. This is shown by Sr and U-Pb isotopic data presented in [14 & 15]. A striking feature of the early Archean Uivak 1 gneisses affected by the late Archean granulite facies event is the obliteration of early Archean fabrics by garnet and orthopyroxene porphyroblasts. This results in the development of a characteristic "blebby texture" [11] in the quartzofeldspathic gneisses. Development of this texture is associated with pervasive migmatization by granitic material. The resulting nebulitic assemblage constitutes the Kiyuktok gneisses [11]. Rocks similar to the Kiyuktok gneisses (albeit, generally strongly deformed) were recognized in west Greenland in 1984 [7]. The Kiyuktok gneisses, together with the Amitsoq gneisses discussed above, are interpreted to represent different evolutionary stages (and perhaps crustal levels) in the chemical and metamorphic differentiation of the gneiss complex in the North Atlantic Craton ca. 2800 to 2900 Ma ago.

Isotopic Data Nd and Sr isotopic data are presented in Figures 1 to 5. The following data presentation for each area is arranged in order of increasing severity of late Archean "reworking".

(1) Type Amitsoq gneisses. Sr isotopic data presented by Moorbath et al., [5] for mixed populations of Amitsoq gneisses yielded ages ranging between 3670 and 3612 Ma (recalculated using  $1.42 \times 10^{-11} \text{ a}^{-1}$  as the  $^{87}\text{Rb}$  decay constant). Initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios ( $I_{\text{Sr}}$ ) for these populations ranged between 0.7001 and 0.7015. Nd isotopic data for samples from the same area are shown in Figure 1. The data show a significant degree of scatter about the reference 3650 Ma isochron (calculated assuming that the  $I_{\text{Nd}}$  had a positive  $\epsilon_{\text{Nd}}$ ). The

Figure 1.



range in  $\epsilon_{\text{Nd}}(0)$  for the samples (ca. -28.5 to -56.4) reflects extreme fractio-

nation in  $^{147}\text{Sm}/^{144}\text{Nd}$  ratios and demonstrates their primitive character.  $T_{\text{CHUR}}^{\text{Nd}}$  model ages range from ca. 3300 to 3680 Ma. Similar Sr and Nd isotopic variations are observed in the best preserved Uivak I gneisses [14,16-19].

(2) Uivak gneisses. Nd isotopic data for the Uivak I gneisses exhibit a range of  $^{147}\text{Sm}/^{144}\text{Nd}$  values between 0.0612 and 0.1197, corresponding to  $\epsilon_{\text{Nd}}(0)$  of -63.1 to -35.4.  $T_{\text{CHUR}}^{\text{Nd}}$  model ages range from ca. 3300 to 3620 Ma. When regressed together with data for the early Archean Nulliak assemblage the total population yields an isochron with a slope equivalent to an age of  $3665 \pm 104$  Ma and an  $I_{\text{Nd}}$  of  $\epsilon_{\text{Nd}} +2.5/-+1$  [14 & 19]. U-Pb ion probe results for Uivak I zircon cores yield Concordia intercept and  $^{207}\text{Pb}/^{206}\text{Pb}$  ages ranging from ca. 3600 to 3900 Ma [2].

(3) Nordafar. Sr isotopic data for a varied suite of 8 samples of Amitsoq gneiss from this area yield a poorly correlated isochron (Fig. 2) equivalent to an age of  $3001 \pm 193/-153$  Ma with an  $I_{\text{Sr}}$  of  $0.7021 \pm 0.0020/+0.0017$  (regressed using




Figure 2




Figure 3




Figure 4




Figure 5

the method of Cameron *et al.*, [17] and assuming partial isotopic homogenization at 1800 Ma). Nd isotopic data for three Nordafar gneisses (Fig. 1) lie within the scatter for the type Amitsoq gneisses. These samples yield  $T_{\text{UR}}^{\text{Sr}}$  model ages between ca. 3400 and 3200 Ma.

(4) Kiyuktok gneisses. Sr and Nd isotopic data for these rocks are discussed in [11, 14, 19-21]. The nebulitic reworked components of the Kiyuktok gneisses comprise varying proportions of old and new components. They have high  $I_{\text{Sr}}$ 's (ca. 0.7055 to 0.7081) and relatively unradiogenic Pb isotopic compositions which reflect involvement of pre-existing crust in their formation. Nd isotopic data for the Kiyuktok gneisses (Fig. 1) plot either within the Uivak I -

Amitsoq gneiss field or above this field, with more enriched radiogenic  $^{143}\text{Nd}$  compositions.

(5) Ikerasakitsup akornga. Sr isotopic data for felsic gneisses from this area are shown in Figure 3. They are moderately well correlated and define an isochron with slope equivalent to an age  $2837 \pm 585/-315$  Ma and an  $I_{\text{Sr}}$  of  $0.7024-21/+12$ . The large uncertainty in age results from limited dispersion in  $^{87}\text{Rb}/^{86}\text{Sr}$  within the suite. Nd isotopic data for specimen (430), the gneiss from this area which gives the oldest  $\text{TUR}^{\text{Sr}}$  model age (3284 Ma), is plotted in Figure 1. The  $\text{TCHUR}^{\text{Nd}}$  model age of this gneiss is 2755 Ma.

(6) Tinissaq. Gneisses from Tinissaq show greater dispersion in their  $^{87}\text{Rb}/^{86}\text{Sr}$  ratios than those from Ikerasakitsup akornga and define an isochron with slope equivalent to an age of  $2858 \pm 214/-163$  Ma and an  $I_{\text{Sr}}$  of  $0.7020-2/+2$  (Fig. 4). When Rb-Sr isotopic data for the Tinissaq and Ikerasakitsup akornga populations are regressed as a regional population using the McIntyre et al., [22] method they define a Model 2 isochron equivalent to an age of  $2906 \pm 136$  Ma with an  $I_{\text{Sr}}$  of  $0.7020-/+3$  (MSWD = 10). Using the Cameron et al., [17] method the age is  $2903 \pm 115/-103$  Ma and the  $I_{\text{Sr}}$  is  $0.7020-/+2$ . Nd isotopic data for a single specimen of felsic gneiss from Tinissaq (425) are shown in Figure 1. This specimen has a slightly more fractionated REE pattern ( $\epsilon_{\text{Nd}}(0) = -31.9$ ) than the specimen from Ikerasakitsup akornga and yields a  $\text{TCHUR}^{\text{Nd}}$  model age of 2821 Ma.


(7) Kangimut sangmissoq. Sr isotopic data for felsic gneisses from this locality (Fig. 5) are poorly correlated and define a McIntyre et al., [22] Model 4 isochron equivalent to an age of  $2768 \pm 264$  Ma with an  $I_{\text{Sr}}$  of  $0.7017-/+5$  (MSWD=79). The Cameron et al., [17] treatment for the same data is also shown in Figure 5. The large uncertainty in the age reflects the influence of the large population with  $^{87}\text{Rb}/^{86}\text{Sr}$  ratios less than 0.1. This depletion in Rb relative to Sr is a characteristic feature of gneisses from this locality. Nd isotopic data for two felsic gneisses are shown in Figure 1. They are less fractionated than the gneisses from the other localities and have  $\epsilon_{\text{Nd}}(0)$  values of -24.6 and -20.5 respectively.  $\text{TCHUR}^{\text{Nd}}$  model ages for these samples (viz., 2625 and 2508 Ma respectively) are significantly lower than model ages displayed by the other felsic gneisses. Also presented in Figure 1 are Sm-Nd isotopic data for a ca. 2900 to 3000 Ma Nuk gneiss from Kangimut sangmissoq, which has an  $\text{TCHUR}^{\text{Nd}}$  model age of 2773 Ma.

Discussion The different groups of felsic gneisses described above are considered to represent different stages in evolution of late Archean crust in the North Atlantic Craton. Sr and Nd isotopic data for most of the gneisses yield equivocal information regarding the crustal prehistory of the gneiss complex. This is particularly true when different groups are studied in isolation. Only the type Amitsoq and Uivak 1 gneisses preserve unequivocal Sr and Nd isotopic evidence of the antiquity of their protoliths. The Kiyuktok gneisses and some of the Nordafar gneisses preserve information about their crustal history in their Sm-Nd isotope systematics (see Fig. 1). However, the Rb-Sr isotopic evolution of the Nordafar gneisses differs from that of the Kiyuktok gneisses and yields no information concerning a possible crustal pre-history (cf. data for the Kiyuktok gneisses, [11 & 21]). This contrasting pattern is interpreted to reflect either variation in metamorphic grade of the early Archean protoliths of these rocks or variable enrichment in Rb during the ca. 2900 Ma metamorphic events (see Nutman et al., [8]). Gneisses at Ikerasakitsup akornga and Tinissaq have similar  $I_{\text{Sr}}$ 's to those at Nordafar (cf. Figs 2 to 4). However, unlike the Nordafar gneisses, their Nd isotope

systematics show no evidence that they contain an older crustal component. Most of the Kangimut sangmissoq gneisses are significantly more depleted in Rb relative to Sr than felsic gneisses from the other areas and are characterized by less highly fractionated REE patterns. Nd isotopic data for these gneisses also show no evidence of an older crustal component.

When combined, the Nd isotopic data for Ikerasakitsup akornga, Tinissaq and Kangimut sangmissoq define a remarkably well correlated array (Fig. 1) with a slope equivalent to an age of ca. 3300 Ma and an extremely depleted  $\epsilon_{Nd}$  (+5.9). On face value this might be interpreted as having "real age significance". However, one of the samples from Kangimut sangmissoq which plots on this line is a ca. 3000 Ma Nuk gneiss also affected by the granulite facies event. This array is therefore interpreted as a pseudoisochron; probably a mixing line without any geochronological significance. The Sm-Nd isotopic systematics of samples which fall on this array, together with the Nordafar and Kiyuktok gneisses, are interpreted to have developed during the ca. 2800 to 2900 Ma tectonothermal event. The data appear to reflect mixing between isotopically evolved Uivak and Amitsoq crust ( $\epsilon_{Nd}$  -8 to -12 at ca. 2900 Ma) and a juvenile-like component, possibly derived from a depleted mantle source with  $\epsilon_{Nd}$  of +4 to +6 at ca. 2900 Ma. The nature of this juvenile component or the mechanism by which the interaction occurs has not yet been established. The model is depicted graphically in Figure 6. Nd isotopic data for Tertiary

Figure 6



basalts flanking, and formed during the opening of Davis Strait, when Labrador was separated from west Greenland clearly have been derived from depleted mantle source regions. The Baffin Bay data, combined with the postulated evolutionary vector for the mantle source region of the early Archean suites, demonstrates that the mantle beneath the North Atlantic Craton would have had the appropriate depleted character at ca. 2900 Ma required by the postulated mixing process. A similar model was suggested by Collerson and McCulloch [14] for the Kiyuktok gneiss and is strengthened by the data presented for the west Greenland gneisses.

## References

- [1] Futa, K. (1981) Sm-Nd systematics of a tonalitic augen gneiss and its constituent minerals from northern Michigan. Geochim. Cosmochim. Acta, 45, 1245-1249.
- [2] Collerson, K.D. (1983) Ion microprobe zircon geochronology of the Uivak gneisses: implications for the evolution of early terrestrial crust in the North Atlantic Craton. In Abstracts for Early Crustal Genesis Field Workshop, 14-18. Lunar and Planetary Institute, Houston.
- [3] McCulloch, M.T. & Black, L.P. (1984) Sm-Nd systematics of Enderby Land granulites and evidence for the redistribution of Sm and Nd during metamorphism. Earth Planet Sci. Lett., 71, 46-58.
- [4] McGregor, V.R. (1973) The early Precambrian gneisses of the Godthaab district, West Greenland. Phil Trans. Roy. Soc. Lond. 273A, 343-358.
- [5] Moorbath, S., O'Nions, R.K., Pankhurst, R.J., Gale, N.H. & McGregor, V.R. (1972) Further rubidium-strontium age determinations on the very early Precambrian rocks of the Godthaab district, west Greenland. Nature Phys. Sci. 240, 78-82.
- [6] Baadsgaard, H. (1973) U-Th-Pb dates on zircons from the Early Precambrian Amitsoq gneisses, Godthaab district, West Greenland. Earth Planet. Sci. Lett. 19, 22-28.
- [7] McGregor et al., (1985) this volume.
- [8] Nutman, A.P., Bridgwater, D. & McGregor, V.R. (1985) Regional variations in the Amitsoq gneisses related to crustal levels during the late Archean granulite facies metamorphism, southern west Greenland. this volume.
- [9] Griffin, W.L., McGregor, V.R., Nutman, A.P., Taylor, P.N. & Bridgwater, D. (1980) Early Archean granulite facies metamorphism south of Ameralik, West Greenland. Earth Planet. Sci. Lett., 50, 59-74.
- [10] Collerson, K.D. & Bridgwater, D. (1979) Metamorphic development of early Archean tonalitic and trondhjemitic gneisses: Saglek area, Labrador. In Trondhjemites, Dacites and Related Rocks (F.Barker, ed.) Elsevier, 205-273.
- [11] \_\_\_\_\_, Kerr, A. & Compston, W. (1981) Geochronology and evolution of late Archean gneisses in Northern Labrador: An example of reworked sialic crust. Spec. Publs. Geol. Soc. Aust., 7, 205-222.
- [12] Ryan, A.B., Martineau, Y., Bridgwater, D., Schiotte, L. & Lewry, J. (1983) The Archean-Proterozoic boundary in the Saglek area, Labrador: Report 1. In Current Research, Part A, Geol. Surv. Can. Pap. 83-1A, 297-304.
- [13] \_\_\_\_\_, \_\_\_\_\_, Korstgaard, J. & Lee, D. (1984) The Archean-Proterozoic boundary in Northern Labrador: Report 2. In Current Research, Part A, Geol. Surv. Can. Pap. 84-1A, 545-551.
- [14] Collerson, K.D. & McCulloch, M.T. (1982) The origin and evolution of Archean crust as inferred from Nd, Sr and Pb isotopic studies in Labrador. 5th Int. Conf. on Geochron. Cosmochron. and Isotope Geol. Extended Abstr. 61-62.
- [15] \_\_\_\_\_, (1983) The Archean gneiss complex of northern Labrador. 2. Mineral ages, secondary isochrons, and diffusion of strontium during polymetamorphism of the Uivak gneisses. Can. J. Earth Sci. 20, 707-718.
- [16] Hurst, R.W., Bridgwater, D., Collerson, K.D. & Wetherill, G.W. (1975) 3600 m.y. Rb-Sr ages from very early Archean gneisses from Saglek Bay, Labrador. Earth Planet. Sci. Lett. 27, 393-403.
- [17] Cameron, M., Collerson, K.D., Compston, W. & Morton, R. (1981) The statistical analysis and interpretation of imperfectly-fitted Rb-Sr isochrons from polymetamorphic terrains. Geochim. Cosmochim. Acta, 45, 1087-1097.
- [18] Collerson, K.D., Brooks, C., Ryan, A.B. & Compston, W. (1982) A reappraisal

- of the Rb-Sr systematics of early Archaean gneisses from Hebron, Labrador. Earth Planet. Sci. Lett. 60, 325-336.
- [19] \_\_\_\_\_, & McCulloch, M.T. (in prep) Sm-Nd systematics of Archaean gneisses from northern Labrador and inferences about crustal evolution.
- [20] \_\_\_\_\_, Kerr, A., Vocke, R.D. & Hanson, G.N. (1982) Reworking of sialic crust as represented in late Archean-age gneisses, northern Labrador. Geology, 10, 202-208.
- [21] Schiotte, L., Bridgwater, D., Nutman, A.P. & Ryan, B. (in press) Effects of late Archaean high-grade metamorphism and granite injection on 3800 Ma gneisses Saglek-Hebron, northern Labrador. Element and isotopic redistribution across an amphibolite-granulite facies transition. J. Geol. Soc.
- [22] McIntyre, G.A., Brooks, C., Compston, W., & Turek, A. 1966. The statistical assessment of Rb-Sr isochrons. J. Geophys. Res. 71, 5459-5468.